

gr-qc/9911062  
UCVFC-DF/16-99

# Analytical Description of Voids in Majumdar-Papapetrou Spacetimes

**Victor Varela<sup>1</sup>**

*Department of Mathematics and Statistics,  
The University of Edinburgh,  
Edinburgh, Scotland*

*Escuela de Física, Facultad de Ciencias,  
Universidad Central de Venezuela,  
Caracas, Venezuela <sup>2</sup>*

E-mail:varela@belka.ciens.ucv.ve

*Departamento de Física,  
Universidad Simón Bolívar,  
Caracas, Venezuela*

## Abstract

We discuss new Majumdar-Papapetrou solutions for the 3+1 Einstein-Maxwell equations, with charged dust acting as the external source of the fields. The solutions satisfy non-linear potential equations which are related to well-known wave equations of 1+1 soliton physics. Although the matter distributions are not localised, they present central structures which may be identified with voids.

---

<sup>1</sup>Talk delivered at the Spanish Relativity Meeting. Bilbao, Spain, September 7-10, 1999.

<sup>2</sup>Present address.

# 1 Introduction

We consider solutions for the Einstein-Maxwell (EM) equations with charged dust acting as the external source of the fields. Our basic equations read

$$G^\mu_\nu = 8\pi T^\mu_\nu, \quad (1)$$

$$F^{\mu\nu}{}_{;\nu} = 4\pi J^\mu, \quad (2)$$

where  $G^\mu_\nu$  and  $F^{\mu\nu}$  denote the Einstein and Maxwell tensors, and the total energy-momentum tensor is given by

$$T^\mu_\nu = E^\mu_\nu + \rho u^\mu u_\nu. \quad (3)$$

Here  $E^\mu_\nu$  is the Maxwell energy-momentum tensor, and the matter term corresponds to dust with energy density  $\rho$  and four-velocity  $u^\mu$ . The four-current is defined by the expression

$$J^\mu = \sigma u^\mu, \quad (4)$$

where  $\sigma$  is the charge density.

We assume that the fluid is static and use the conformal static metric

$$ds^2 = -V^2 dt^2 + \frac{1}{V^2} h_{ij} dx^i dx^j, \quad (5)$$

where the background metric  $h_{ij}$  and  $V$  depend only on the space-like coordinates  $x^1, x^2, x^3$ . The electrostatic forms of  $A_\mu$  and  $J^\mu$  are given by

$$A_\mu = A_0(x^i) \delta_\mu^0, \quad (6)$$

$$J^\mu = \frac{\sigma(x^i)}{V} \delta_0^\mu, \quad (7)$$

with  $i = 1, 2, 3$ .

Under these conditions, Eq. (2) contains only one non-trivial equation:

$$\frac{1}{\sqrt{h}} \partial_j \left( \sqrt{h} h^{jk} \frac{\partial_k A_0}{V^2} \right) = \frac{4\pi J^0}{V^2}, \quad (8)$$

where  $h$  and  $h^{ij}$  are the determinant and the inverse of  $h_{ij}$ , respectively.

The trace of the Einstein equations is

$$R = -8\pi T, \quad (9)$$

where  $R$  denotes the Ricci scalar and  $T = T^\mu_\mu$ . We use the decomposition

$$R = V^2 \left[ R_h + 2\nabla_h^2 \ln V - 2\partial_i \ln V \partial^i \ln V \right]. \quad (10)$$

Here  $R_h$  is the Ricci scalar associated to  $h_{ij}$ , and  $\nabla_h^2$  is the three-dimensional Laplacian operator constructed with the same metric. We assume a flat background space, with  $R_h = 0$ . Therefore, combining Eqs. (9) and (10) we obtain

$$\nabla_h^2 \left( \frac{1}{V} \right) = \frac{4\pi T}{V^3}. \quad (11)$$

Following the Majumdar-Papapetrou (MP) procedure,[1, 2] we assume that

$$A_0 = \alpha V, \quad (12)$$

where  $\alpha = \pm 1$ . As a consequence, the Maxwell equation (8) takes the form

$$\nabla_h^2 \left( \frac{1}{V} \right) = -\frac{4\pi\alpha J^0}{V^2}, \quad (13)$$

which is clearly the same as Eq. (11) whenever the condition

$$T = -\alpha J^0 V \quad (14)$$

holds. This equation can be combined with  $J^0 = \frac{\sigma}{V}$  to obtain the alternative expression

$$\sigma = -\alpha T. \quad (15)$$

Since  $T = -\rho$  for dust, Eqs. (11) and (15) can be finally expressed as

$$\nabla_h^2 \lambda + 4\pi\rho\lambda^3 = 0, \quad (16)$$

$$\sigma = \alpha\rho, \quad (17)$$

where  $\lambda = \frac{1}{V}$ . Due to Eqs. (5) and (12), only one Einstein equation is not trivially satisfied.[3, 4] Therefore, solving Eq. (16) is sufficient for finding a solution of the EM equations.

If we identify our flat background space with the Euclidean, three dimensional space and assume  $\rho = 0$ , then Eq. (16) reduces to the usual Laplace equation  $\nabla^2 \lambda = 0$  and the electrovac, multi-black hole solution follows straightforwardly. Assuming spherical symmetry, and using spherical coordinates, we find

$$\lambda = 1 + \frac{m}{r}. \quad (18)$$

In the far-asymptotic region, the behaviour of this solution is approximately given by

$$V \approx 1 - \frac{m}{r}, \quad g_{00} \approx -1 + \frac{2m}{r}, \quad A_0 \approx \pm(1 - \frac{m}{r}). \quad (19)$$

The corresponding expression for the electric field is

$$E \approx \frac{q}{r^2}, \quad (20)$$

where

$$q = \pm m. \quad (21)$$

Equation (5) implies that the invariant area of any 2-sphere surrounding the origin is given by  $\frac{4\pi r^2}{V(r)^2}$ . Therefore, the set  $r = 0$ ,  $t = \text{constant}$  has a non-zero invariant area given by  $4\pi m^2$ . In fact, a simple coordinate transform shows that the null hypersurface  $r = 0$  is the horizon of the extremal Reissner-Nordström solution. Also, if we define the new radial coordinate  $\tilde{r} = -r$  and perform the standard analysis,[5] then we find

that this horizon encloses a point-like, essential singularity placed at  $\tilde{r} = m$ . In fact, the invariant area vanishes and the scalar  $J = F_{\mu\nu}F^{\mu\nu} = \lambda^{-4} \left(\frac{d\lambda}{dr}\right)^2$  blows up at that point.

Equations (16) and (17) were originally discussed by Das [6] in his study of equilibrium configurations of self-gravitating, charged dust. More recently, Gürses [3] has considered non-electrovac solutions when Eq. (16) is linear. This situation corresponds to his choice  $\rho = \frac{b^2}{4\pi\lambda^2}$  for constant  $b$ . In this case, Eq. (16) admits the particular solution  $\lambda = \frac{a \sin br}{r}$  where  $a$  is an integration constant. The oscillatory behaviour of this solution implies a geometry with a complicated radial dependence. In fact, the invariant area vanishes for a discrete, infinite set of values of  $r$ , and the Ricci scalar  $R = \frac{2b^2 r^2}{a^2 \sin^2 br}$  blows up wherever the invariant area vanishes, except for  $r = 0$ . Other solutions with oscillatory behaviour have been considered by Balakrishna and Wali,[7] Braden and Varela,[8] and Ida.[4] In Section 2 we exploit the general non-linearity of Eq. (16) to obtain new solutions which are free of oscillatory singularities and allow asymptotically flat behaviour.

## 2 The non-linear models

The non-linear potential equation (16) takes the spherically symmetric form

$$\frac{d^2\lambda}{dr^2} + \frac{2}{r} \frac{d\lambda}{dr} + 4\pi\rho\lambda^3 = 0. \quad (22)$$

Using the new radial coordinate  $\tau = \frac{1}{r}$ , the same differential equation can be written as

$$\frac{d^2\lambda}{d\tau^2} + \frac{4\pi\rho}{\tau^4}\lambda^3 = 0. \quad (23)$$

If  $\rho$  and  $\lambda$  satisfy the condition

$$\rho = \frac{b^2}{4\pi} \frac{\tau^4 \sin \lambda}{\lambda^3}, \quad (24)$$

then (23) finally reduces to the -sine-Gordon equation [9]

$$\frac{d^2\lambda}{d\tau^2} + b^2 \sin \lambda = 0, \quad (25)$$

which has the solutions

$$\lambda^\pm(\tau) = 2 \arcsin [\tanh(\pm b\tau + c)] + 2n\pi, \quad (26)$$

where  $n$  is an arbitrary integer,  $c$  is an integration constant, and  $b$  is assumed to be positive. We consider only the case  $n = 0$ . In terms of the original radial coordinate, these solutions read

$$V^\pm(r) = \frac{1}{2 \arcsin \left[ \tanh \left( \pm \frac{b}{r} + c \right) \right]}. \quad (27)$$

We observe that  $V^\pm(0)^2$  is finite, so the invariant area vanishes for  $r = 0$ . Therefore, the set  $r = 0, t = \text{constant}$  is point-like with respect to both solutions. Let us deal with  $V^+$  first. A preliminary numerical study of the invariants  $J, R, R^{\alpha\beta}R_{\alpha\beta}, R^{\alpha\beta\gamma\delta}R_{\alpha\beta\gamma\delta}$  suggests that these quantities are bounded for non-negative  $r$ , whenever  $c$  is positive. If we choose

$$c = \frac{1}{2} \ln \left[ \frac{1 + \sin(1/2)}{1 - \sin(1/2)} \right], \quad (28)$$

then the far-asymptotic behaviour of this solution is given by Eqs. (19), (20), (21) with  $m = 2b \cos(1/2)$ . Therefore,  $V^+$  is asymptotically flat, exactly as the MP electrovac solution.

The positive definite energy density given by Eq. (24) corresponds to a non-localised matter (and charge) distribution. However,  $\rho$  is negligible for  $x = \frac{r}{b} \ll 0.2$ . For very small  $x$  the dimensionless expressions of  $\rho$  and  $R^{\alpha\beta\gamma\delta}R_{\alpha\beta\gamma\delta}$  are approximately given by

$$\rho(x) \approx \frac{e^{-c}}{\pi^4} \frac{e^{-\frac{1}{x}}}{x^4}, \quad (29)$$

$$R^{\alpha\beta\gamma\delta}R_{\alpha\beta\gamma\delta}(x) \approx 3 \left( \frac{2}{\pi} \right)^6 e^{-2c} \frac{e^{-\frac{2}{x}}}{x^8}. \quad (30)$$

These results imply a very fast decrease of  $\rho(x)$  and  $R^{\alpha\beta\gamma\delta}R_{\alpha\beta\gamma\delta}(x)$  when  $x \rightarrow 0^+$ , and suggest the existence of a void in the innermost region of the asymptotically flat object constructed with  $V^+$ . Nevertheless, this interpretation cannot be complete without a better understanding of the point-like set  $r = 0, t = \text{constant}$ . A closer look at the singularity contained in this solution is also necessary. We interpret  $r = 0, t = \text{constant}$  as the center of symmetry and observe that the above mentioned invariants are bounded at this point. However, the coordinate transform  $\tilde{r} = -r$  reveals the existence of a point-like, essential singularity at  $\tilde{r} = \frac{b}{c}$ . In fact, the invariant  $J$  blows up at this point. The use of  $V^+$  alone may imply the division of the manifold into connected parts, separated by the point-like singularity placed at  $r = -\frac{b}{c}$ . However, a very different situation comes out when we restrict  $V^+$  to positive values of  $r$  and describe the geometry for  $r < 0$  with the second solution  $V^-$ . Then, a smooth (at least  $C^1$ ) matching of  $V^+$  and  $V^-$  occurs at  $r = 0$  and the arising asymptotically flat spacetime seems to be connected and singularity free, with an almost empty region near the center of symmetry. Thus, the joint use of  $V^+$  and  $V^-$  provides a simpler description of a MP void. The study of the global structure of these solutions is left as an open problem, which provides motivation for further research work.

Finally, we point out that other exact, non-linear solutions for this theory can be found if we impose different relationships between  $\rho$  and  $\lambda$ . For example, the choice

$$\rho = -\frac{b^2}{4\pi} \frac{\tau^4 \sin \lambda}{\lambda^3} \quad (31)$$

leads to the sine-Gordon equation

$$\frac{d^2 \lambda}{d\tau^2} = b^2 \sin \lambda. \quad (32)$$

It has the well-known solutions

$$\lambda^\pm(\tau) = 4 \arctan e^{(\pm b\tau + d)}. \quad (33)$$

If we choose  $d = \ln[\tan(1/4)]$ , then both solutions have asymptotically flat behaviour.

Another example is

$$\rho = \frac{b^2}{4\pi} (\lambda - \lambda^3). \quad (34)$$

In this case the geometry is determined by the  $\lambda\phi^4$  equation

$$\frac{d^2\lambda}{d\tau^2} + b^2 (\lambda - \lambda^3) = 0 \quad (35)$$

which admits the solutions

$$\lambda^\pm(\tau) = \tanh\left(\pm \frac{b}{\sqrt{2}}\tau + f\right). \quad (36)$$

The relationship between the 3+1 EM theory and the equations of 1+1 soliton physics deserves a more detailed examination. Possible extensions of this work involve the analysis of dust models for which  $\lambda(\tau)$  is a solution of the KdV equation, and the study of the non-linear potential equations arising in higher dimensions.[10]

## Acknowledgments

The author is grateful to the Mathematics and Statistics Department of Edinburgh University for its hospitality during his long-term research visit. He is indebted to Professors Nikolaos Batakis (Ioannina) and Luis Herrera (Salamanca) for useful discussions and comments. Finally, he thanks Dr. Pio J. Arias (UCV) and Professor Jorge Zanelli (Santiago) for valuable suggestions, and the *Grupo de Relatividad y Campos* at Simón Bolívar University for partial financial support.

## References

- [1] S. D. Majumdar, Phys. Rev., **72**, 390 (1947).
- [2] A. Papapetrou, Proc. Roy. Irish Academy A, **51**, 191 (1947).
- [3] M. Gürses, Phys. Rev. D, **58**, 044001 (1998).
- [4] D. Ida, gr-qc/9906071.
- [5] J. B. Hartle and S. W. Hawking, Commun. Math. Phys., **26**, 87 (1972).
- [6] A. Das, Proc. R. Soc. London A, **267**, 1 (1962).
- [7] B. S. Balakrishna and K. C. Wali, Phys. Rev. D, **46**, R5228 (1992).
- [8] H. W. Braden and V. Varela, Phys. Rev. D, **58**, 124020 (1998).
- [9] B. Saha (gr-qc/9811044) has considered scalar fields satisfying this and other solitonic equations on a static Freedman-Robertson-Walker background.
- [10] The electrovac MP solutions in higher dimensions have been discussed by P. Mitra (gr-qc/9908014).